

## Exhibit 8

# Photochemistry and Photobiology

---

~~Contents, Author and Subject Index~~

Volume 68, 1998

Published by  
the American Society  
for Photobiology

# Photochemistry and Photobiology

The official journal of the American Society for Photobiology

## Editor

J. C. Scaiano, Ph.D.  
Department of Chemistry  
University of Ottawa  
10 Marie Curie, Rm. 306  
Ottawa, ON K1N 6N5 Canada  
Phone: 613-562-5634  
Fax: 613-562-5633  
e-mail: P&P@photo.chem.uottawa.ca

## Associate Editors

Jean Cadet, *CEA/Département de Recherche  
Fondamentale sur la Matière Condensée,  
Grenoble, France*

Frank R. de Gruijl, *Utrecht University  
The Netherlands*

Xing Wang Deng, *Yale University  
New Haven, CT*

J. Woodland Hastings, *Harvard University,  
Cambridge, MA*

Barbara W. Henderson, *Roswell Park  
Cancer Institute, Buffalo, NY*

Margaret L. Kripke, *Anderson Cancer  
Center, Houston, TX*

Miguel Angel Miranda, *Universidad  
Politecnica de Valencia, Valencia, Spain*

Thomas A. Moore, *Arizona State  
University, Tempe, AZ*

Harald Paulsen, *Johannes-Gutenberg-  
Universität, Mainz, Germany*

Aziz Sancar, *The University of North  
Carolina at Chapel Hill, Chapel Hill, NC*

Gary B. Schuster, *Georgia Institute of  
Technology, Atlanta, GA*

Eva M. Sevcik-Muraca, *Purdue University,  
West Lafayette, IN*

John L. Spudich, *University of Texas  
Medical School, Houston, TX*

Manfred Tevini, *Universität Karlsruhe,  
Karlsruhe, Germany*

Dennis P. Valenzano, *University of Kansas  
Medical Center, Kansas City, KS  
(electronic publication)*

Brian C. Wilson, *Ontario Cancer Institute,  
Toronto, ON*

**Photochemistry and Photobiology** (ISSN 0031-8655) is published monthly, two volumes per annum, by the American Society for Photobiology, BioTech Park, 1021 15th Street, Suite 9, Augusta, GA 30901-3158, USA.

**Subscriptions:** The 1998 Annual Institutional Subscription Rate is \$575 for 12 monthly issues plus Meeting Abstract issue. Institutional subscription information is available from Allen Press, Inc., 1041 New Hampshire Street, P.O. Box 368, Lawrence, KS 66044; Phone 785-843-1221, Fax 785-843-1274. Individual members of the American Society for Photobiology receive a complimentary subscription. Membership information may be obtained from the Society office at the address listed above or at <http://www.POL-US.net>.

**Instructions to Authors:** Condensed instructions appear at the end of each issue. Complete instructions for preparation of manuscripts and figures and for submission of manuscripts on disk appear in the first issue of each year and are available from the Editorial Office: *Photochemistry and Photobiology*, Department of Chemistry, University of Ottawa, 10 Marie Curie, Rm. 306, Ottawa, ON K1N 6N5 Canada. Phone: 613-562-5634, Fax: 613-562-5633, e-mail: P&P@photo.chem.uottawa.ca. Complete information is also available on the *Photochemistry and Photobiology* home page.

**Internet address:** The address for the *Photochemistry and Photobiology* home page is <http://www.POL-US.net>. The home page provides information on published papers and manuscripts prior to publication. It also includes information on subscriptions, the aims and scope of the Journal, preparation of manuscripts, and advertisements.

**Advertisements:** Advertising rates and information are available from the Editorial Office at the address listed above and on the home page.

**Change of address:** Institutional subscribers should forward their new address to Allen Press at the address for subscriptions given above. Members of the American Society for Photobiology should send new addresses to the Society office listed above.

**Back issues:** Claims for undelivered issues must be received within six months of the mailing date. Single issues and complete volumes can be purchased depending on availability. Contact Allen Press at the address for subscriptions given above.

**Indexing/Abstracting Services:** The Journal is currently included by the following services in print and/or electronic format: Current Contents (Life Sciences, Science Citation Index, SCISEARCH Data, ISI/BIOMED Database, ASCA), Chemical Abstracts Service, Index Medicus, MED-LINE, BIOSIS Data, Cambridge Science Abstracts, CABS, and PASCAL-CNRS Data.

**Copyright information:** *Photochemistry and Photobiology* is copyrighted by the American Society for Photobiology. No portion(s) of this journal may be reproduced without the written consent from the copyright holder. Permission to reproduce copies of articles for non-commercial use may be obtained from the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923, 508-750-8400, for a fee of \$5.00 per copy.

**Periodicals postage paid at Augusta, GA and additional mailing offices.**

**Postmaster:** Send address corrections to *Photochemistry and Photobiology*, American Society for Photobiology, 1021 15th Street, Suite 9, Augusta, GA 30901-3158.

**Editorial Advisory Board:** P. L. Dutton, Philadelphia, PA; J. E. Hearst, Berkeley, CA; L. H. Pratt, Athens, GA; R. B. Setlow, Upton, NY; B. C. Wilson, Toronto, Canada

**Founding Editors:** E. J. Bowen, Oxford, UK; S. Claesson, Uppsala, Sweden; A. Hollaender, Oak Ridge, TN; D. Shugar, Warsaw, Poland; A. D. McLaren, Berkeley, CA (Editor, 1962-1965)

**Past Editors:** K. C. Smith, Stanford, CA (1966-1971); J. Jagger, Dallas, TX (1972-1974); P.-S. Song, Lincoln, NE (1975-1993); Irene E. Kochevar, Boston, MA (1994-1997)

**Editorial Associate:** Diann H. Denning

**Editorial Assistants:** Anita Bouwman and Adriana Laliberté

Copyright © 1998 American Society for Photobiology

The paper used in this publication meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of paper)

## Preferential Relative Porphyrin Enrichment in Solar Keratoses upon Topical Application of $\delta$ -Aminolevulinic Acid Methyl ester

Clemens Fritsch<sup>\*1,2</sup>, Bernhard Homey<sup>2</sup>, Wilhelm Stahl<sup>1</sup>, Percy Lehmann<sup>2</sup>, Thomas Ruzicka<sup>2</sup> and Helmut Sies<sup>1</sup>

<sup>1</sup>Institute of Physiological Chemistry I and <sup>2</sup>Department of Dermatology, Heinrich-Heine-University, Düsseldorf, Germany

Received 5 January 1998; accepted 15 May 1998

### ABSTRACT

Topically applied  $\delta$ -aminolevulinic acid is used efficiently for the treatment of solar keratoses by photodynamic therapy. Recent animal studies suggest that porphyrin sensitization of epithelial tissue is improved by using esters rather than free  $\delta$ -aminolevulinic acid. The present study examines porphyrin metabolite formation after topical application of  $\delta$ -aminolevulinic acid or  $\delta$ -aminolevulinic acid methyl ester in human solar keratoses *versus* adjacent normal skin. Levels of total porphyrins, porphyrin metabolites and protein were measured in skin samples excised after 1 and 6 h. Higher levels of porphyrins were observed in solar keratoses than in normal skin with both substances. Maximum porphyrin levels were present in solar keratoses treated with  $\delta$ -aminolevulinic acid for 6 h. However, the ratio of porphyrins in solar keratoses *versus* adjacent normal skin was higher with  $\delta$ -aminolevulinic acid methyl ester. The pattern of porphyrins showed no significant difference between normal and afflicted skin, protoporphyrin being predominant. The results suggest that application of free  $\delta$ -aminolevulinic acid may be more effective in sensitizing solar keratoses. However, treatment with  $\delta$ -aminolevulinic acid methyl ester leads to a preferential enrichment of porphyrins within lesional skin.

### INTRODUCTION

Photodynamic therapy (PDT)<sup>†</sup> with  $\delta$ -aminolevulinic acid hydrochloride (ALA) is based on the administration of ALA to the diseased skin and subsequent irradiation of the porphyrin-sensitized tissue with red light (1). Tumor necrosis is induced by reactive oxygen species targeting biological membranes as well as by vascular damage leading to tumor cell anoxia (2). Topical ALA-PDT was shown to be highly effective in the treatment of solar keratoses (SK), superficial

basal cell carcinoma and superficial squamous cell carcinoma (1,3-6) without serious adverse effects (7).

Accumulation of high intralosomal porphyrin levels is essential for effective ALA-PDT. Previously, we showed that irradiation 1-6 h after ALA application was optimal in superficial epithelial skin tumors in relation to adjacent normal skin (8). However, the homogeneity of porphyrin sensitization of neoplastic cutaneous tissues needs to be augmented for improved response rates in ALA-PDT. Higher lipophilicity of ALA esters permits more effective penetration of cutaneous tissue as compared to free ALA. In a recent study, the topical application of ALA esters led to higher porphyrin fluorescence in mouse skin when compared with ALA hydrochloride (9).

Therefore, here we examined the formation of porphyrins after topical application of ALA and ALA methyl ester (ALA-ME) in SK and adjacent normal skin (NS).

### MATERIALS AND METHODS

**Materials.** Tissue samples of SK and NS adjacent to the lesions (each  $n = 40$ ) were obtained from patients who underwent surgery. Tissue samples were taken as follows: without substrate application ( $n = 8$ ), 1 h after application of ALA ( $n = 8$ ) or ALA-ME ( $n = 8$ ) and 6 h after application of ALA ( $n = 8$ ) or ALA-ME ( $n = 8$ ). One part of the excised tissue was examined histopathologically. Each patient received comprehensive information about the scope of the study.

**ALA treatment and photodynamic diagnosis (PDD).** Based on pilot studies of topical ALA application (8) and on clinical experience in ALA-PDT (3,4,6), of 20% ALA hydrochloride (Merck, Darmstadt Germany) or the short-chain methyl ester (Sigma Aldrich Chemie, Deisenhofen, Germany) were mixed in an ointment (Neribas<sup>®</sup>, Schering, Berlin, Germany) and 0.2 g of this mixture was applied to a 1 cm<sup>2</sup> skin area ( $\approx 40$  mg/cm<sup>2</sup>) of SK or NS. Treated skin was covered with an occlusive foil (Tegaderm<sup>®</sup>, 3M Healthcare, Borken, Germany), gauze, aluminum foil and tape to enhance tissue penetration and avoid photobleaching of porphyrins. After 1 or 6 h, the ointment was removed and the treated area was illuminated with Wood's light (Fluotest<sup>®</sup>, Xenotest, Hanau, Germany; 370-405 nm). Detected fluorescence intensity was expressed semiquantitatively relative to a fluorescence standard. The fluorescent area was marked. Basal values were obtained from untreated controls.

**Preparation of skin samples.** Only superficial layers of skin ( $<1$  mm) were included in the study because of the limited penetration of topically applied ALA (10-12). Immediately after excision, tissue samples were frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$ .

**Determination of total porphyrin and protein levels.** Tissue samples were weighed and cut into small pieces. After homogenization with an Ultraturrax and centrifugation at 3000 U/min for 10 min, porphyrins were isolated with 1.0 N perchloric acid/methanol (1/1,

\*To whom correspondence should be addressed at: Institut für Physiologische Chemie, Heinrich-Heine-Universität Düsseldorf, 40001 Düsseldorf, Germany. Fax: 49-211-81-13029.

†Abbreviations: ALA,  $\delta$ -aminolevulinic acid; ALA-ME,  $\delta$ -aminolevulinic acid methyl ester; NS, normal skin; PDD, photodynamic diagnosis; PDT, photodynamic therapy; SK, solar keratoses.

© 1998 American Society for Photobiology 0031-8655/98 \$5.00+0.00

**Table 1.** Fluorescence intensities of tissues treated by  $\delta$ -aminolevulinic acid or  $\delta$ -aminolevulinic acid methylester\*

	$\delta$ -Aminolevulinic acid		$\delta$ -Aminolevulinic acid methylester	
	1 h	6 h	1 h	6 h
Normal skin	1.1 $\pm$ 0.2	1.9 $\pm$ 0.3	0.5 $\pm$ 0.2	0.8 $\pm$ 0.2
Solar keratoses	1.9 $\pm$ 0.2	2.9 $\pm$ 0.1	1.4 $\pm$ 0.2	1.9 $\pm$ 0.2

\*Fluorescence intensities of skin treated by  $\delta$ -aminolevulinic acid or  $\delta$ -aminolevulinic acid methylester irradiated by Wood's light (370–405 nm, 10 cm distance, 5 mW/cm<sup>2</sup>). Values are given according to a fluorescence standard: 0 = no fluorescence, 3 = maximum fluorescence ( $n = 8$ ; mean  $\pm$  SEM).

vol/vol). In the supernatant, the total porphyrin level was assessed by fluorescence spectroscopy (Perkin Elmer LS-5, Überlingen, Germany); emission was recorded between 520 and 700 nm at an excitation wavelength of 405 nm (Soret band). A protoporphyrin standard was obtained from Porphyrin Products (Logan, UT, USA) (13). Protein levels were determined in the pellet (14).

The relative porphyrin enrichment was expressed by the ratio of porphyrin accumulation in SK versus adjacent NS.

**Porphyrin metabolites.** The supernatant was adjusted with acetic acid to pH 3–4, porphyrins were bound to talcum, esterified and metabolites were identified by HPLC with fluorescence detection (L-7480, Merck Hitachi, Darmstadt Germany) using a porphyrin standard mixture (Porphyrin Products) for quantification (15,16). The following metabolites were analyzed: protoporphyrin, tricarboxylic porphyrin, coproporphyrin, pentacarboxylic porphyrin, hexacarboxylic porphyrin, heptacarboxylic porphyrin and uroporphyrin. The latter four porphyrins are assigned as highly carboxylated porphyrins.

**Statistical calculation.** Statistical analysis was performed by Student's *t*-test. Data are reported as means  $\pm$  SEM. Changes were considered statistically significant when the *P* value was  $<0.05$ .

## RESULTS

### Fluorescence intensities of treated tissues

Fluorescence intensity was higher in SK topically treated with ALA for 6 h. Fluorescence was weaker for ALA-ME than for ALA in normal skin (Table 1).


### Basal total porphyrin levels

In untreated tissues, the levels of total porphyrins were similarly low: 1.2  $\pm$  0.1 in normal skin and 1.3  $\pm$  0.1 nmol/g protein in SK (Fig. 1 and Table 2).

### Total porphyrin levels after topical application of ALA or ALA-ME

Topical application of both compounds led to increased porphyrin levels in SK when compared with NS. Highest levels were detected at 6 h. The ALA treatment induced highest porphyrin values: 36.4  $\pm$  4.0 in SK and 7.2  $\pm$  0.5 nmol/g protein in NS adjacent to the lesions (Fig. 1A). Using ALA-ME as substrate, porphyrin levels in SK were 14.9  $\pm$  2.2 nmol/g protein, which is less than 50% of the amount induced by ALA (Fig. 1B). Furthermore, lowest porphyrin levels were detected in NS treated with ALA-ME (1.7  $\pm$  0.2 nmol/g protein), which is four times less than for ALA.

The relative porphyrin enrichment, i.e. the ratio of the level in SK versus NS (SK/NS), showed maximum values at 6 h: 5.1 for ALA and 8.7 for ALA-ME (Fig. 2).



**Figure 1.** Porphyrin levels in normal skin (NS, open squares) and solar keratoses (SK, filled squares) without and 1 and 6 h after topical application of  $\delta$ -aminolevulinic acid (ALA) or  $\delta$ -aminolevulinic acid methylester (ALA-ME). A. Using ALA, in both tissues, maximum porphyrin levels were measured at 6 h. The SK accumulated more porphyrins (36.4  $\pm$  4.0) than NS (7.2  $\pm$  0.5 nmol/g protein; \**P* < 0.005 SK vs NS). B. Using ALA-ME, porphyrin accumulation was less than with free ALA (A). In SK increased porphyrin synthesis was detected when compared with NS. Distinct differences already occurred at 1 h: SK 3.7  $\pm$  0.5 and NS 1.3  $\pm$  0.1 (\**P* < 0.05 SK vs NS). Maximum porphyrin levels were measured at 6 h (SK: 14.9  $\pm$  2.2 and NS: 1.7  $\pm$  0.2; \*\**P* < 0.01 SK vs NS).

### Porphyrin metabolite levels

The basal pattern of porphyrin metabolites was comparable in both untreated tissues with protoporphyrin as the predominant metabolite (89–91%) followed by uroporphyrin and coproporphyrin (Table 2). The pattern of metabolites remained unchanged upon treatment. Protoporphyrin was still the predominant metabolite (82–91%) and accumulated in both tissues over the period of 6 h (Table 2).

## DISCUSSION

Porphyrin metabolite formation in human SK and adjacent NS after topical application of ALA-ME versus ALA was investigated in the present study. Highest levels of porphyrins are achieved after 6 h in both tissues. Recent biochemical studies on the time course of ALA-induced porphyrin accumulation in epithelial skin tumors showed that maximum intralesional porphyrin levels were measured between 1 and 6 h (8). Therefore, the present study was focused on the application times 1 and 6 h.

Porphyrin accumulation was more intense using ALA than ALA-ME, suggesting that ALA may penetrate the epidermal barrier with subsequent conversion into porphyrins slightly

Table 2. Porphyrin levels and distribution of porphyrin metabolites 6 h after substrate application\*

Substrate	Total porphyrins (nmol/g protein)	Proto-porphyrin (%)	Copro-porphyrin (%)	Pentacarboxylic porphyrin (%)	Hexacarboxylic porphyrin (%)	Heptacarboxylic porphyrin (%)	Uroporphyrin (%)
Normal skin							
No	1.2 ± 0.1	91 ± 7	3 ± 1	1 ± 1	0 ± 1	0 ± 4	1 ± 3
ALA	7.2 ± 0.5	90 ± 2	0 ± 1	0 ± 0	0 ± 0	4 ± 1	7 ± 1
ALA-ME	1.7 ± 0.2	89 ± 2	0 ± 0	0 ± 0	1 ± 0	4 ± 0	6 ± 1
Solar keratoses							
No	1.3 ± 0.1	89 ± 7	3 ± 3	0 ± 0	0 ± 1	1 ± 1	7 ± 9
ALA	36.4 ± 4.0*	90 ± 1	1 ± 1	0 ± 0	2 ± 1	2 ± 0	3 ± 1
ALA-ME	14.9 ± 2.2†	82 ± 4	8 ± 2	0 ± 1	2 ± 1	5 ± 1	5 ± 1

\* Data are given as nmol porphyrin/g protein and percent total porphyrins ( $n = 8$ ; mean ± SEM; \* $P < 0.005$ , † $P < 0.01$ : solar keratoses vs normal skin). Porphyrin patterns showed no significant differences between solar keratoses and normal skin. ALA =  $\delta$ -aminolevulinic acid; ALA-ME =  $\delta$ -aminolevulinic acid methylester.

more efficiently than ALA-ME, which requires hydrolysis after penetration. However, using ALA the lesion-adjacent normal skin also accumulated relatively high porphyrin amounts. Thus, topical application of ALA-ME diminishes porphyrin sensitization in normal skin. Regarding the relative enrichment in porphyrin levels SK/NS, ALA-ME seems preferentially to penetrate the damaged skin and abnormal cells of SK (Fig. 2). The low levels of porphyrins in perilesional NS might be primarily due to weak penetration of ALA-ME through intact skin, less formation of porphyrins out of the ester form due to slower deesterification in normal keratinocytes or lower leakage of intralesional formed porphyrins into the perilesional tissue as previously postulated for ALA (8).

The formation of porphyrins after topical application of ALA esters was recently investigated in murine skin (9). All ALA derivatives studied there (methylester, ethylester and propylester) showed higher induction of porphyrin fluorescence than free ALA, particularly after a prolonged application time of 14 h. In contrast, our human *in vivo* data show that NS is less sensitized with porphyrins using ALA-ME when compared with ALA. The use of different species and the lack of neoplastic tissue in the mice might explain the different results. It is known that perilesional NS, as exam-

ined in our study, accumulates about three times more porphyrin than NS distant from lesions such as squamous cell carcinomas when treated topically by ALA (8). Thus, it is surprising that the NS localized close to the SK accumulated only low porphyrin levels upon treatment with ALA-ME. Specificity of ALA-ME versus ALA may also differ depending on the time of application.

The porphyrin metabolite pattern shows that any disturbance of the heme-biosynthesis-associated enzymes seems to be not relevant in human SK. The accumulation of higher total porphyrin levels appears to be the major effect responsible for intralesional photosensitization.

The penetration depth of porphyrin fluorescence induced by topical ALA was shown to be inhomogenous (10–12,17–19). Further studies have to show whether ALA esters are superior to free ALA with respect to homogenous porphyrin formation in neoplastic lesions.

In topical ALA-PDT, local photosensitivity commonly lasts up to 48 h (6). The higher lesional selectivity of ALA-ME-induced porphyrins provides for a reduction in cutaneous light sensitivity, particularly in perilesional NS (8). Based on *in vitro* data it might be speculated that long-chain ALA esters are more efficient in PDT than ALA or short-chain ALA esters (20). Optimum efficacy of topical PDT may be expected if irradiation of SK is performed 6 h after treatment with ALA-ME. Application of ALA-ME also optimizes photodynamic diagnosis (PDD), a novel effective diagnostic modality in tumor detection and demarcation (5,6,21,22).

Therefore, the use of ALA-esters in PDT and PDD seems to be a promising modality with highly selective intralesional porphyrin formation.

**Acknowledgement**—C.F. has been supported by a grant of the Deutsche Forschungsgemeinschaft (Fr 1174/1–1).

## REFERENCES

- Kennedy, J. C., R. H. Pottier and D. C. Pross (1990) Photodynamic therapy with endogenous protoporphyrin IX: basic principles and present clinical experience. *J. Photochem. Photobiol. B Biol.* 6, 143–148.
- Pass, H. I. (1993) Photodynamic therapy in oncology: mechanisms and clinical use. *J. Natl. Cancer Inst.* 85, 443–453.
- Wolf, P., E. Rieger and H. Kerl (1993) Topical photodynamic therapy with endogenous porphyrins after application of  $\delta$ -aminolevulinic acid. *J. Am. Acad. Dermatol.* 28, 17–21.




Figure 2. Ratio of total porphyrins in solar keratoses (SK) versus normal skin (NS) after topical application of 20%  $\delta$ -aminolevulinic acid (ALA; open squares) or  $\delta$ -aminolevulinic acid methylester (ALA-ME; filled squares). The ALA-ME treatment produced significantly higher ratios of porphyrin levels (SK/NS) when compared with ALA treatment: at 1 h the ratio was 2.9 (\* $P < 0.05$  ALA-ME vs ALA) and at 6 h 8.7 (\*\* $P < 0.01$  SK vs NS) was measured.

4. Calzavara-Pinton, P. G. (1995) Repetitive photodynamic therapy with topical 5-aminolevulinic acid as an appropriate approach to the routine treatment of superficial non-melanoma skin tumours. *J. Photochem. Photobiol. B Biol.* **29**, 53-57.
5. Fritsch, C., P. M. Becker-Wegerich, K. W. Schulte, W. Neuse, P. Lehmann, T. Ruzicka and G. Goerz (1996) Treatment of a large superficial basal cell carcinoma of the breast with combination of photodynamic therapy and surgery controlled by photodynamic diagnosis. *Hautarzt* **47**, 438-442.
6. Fritsch, C., G. Goerz and T. Ruzicka (1998) Photodynamic therapy in dermatology. A review. *Arch. Dermatol.* **134**, 207-214.
7. Fritsch, C., B. Verwohlt, K. Bolsen, T. Ruzicka and G. Goerz (1996) Influence of topical photodynamic therapy with 5-aminolevulinic acid on the porphyrin metabolism. *Arch. Dermatol. Res.* **288**, 517-521.
8. Fritsch, C., W. Stahl, K. W. Schulte, H. Sies, G. Goerz, L. P. Lehmann and T. Ruzicka (1998) Optimum porphyrin formation in skin tumors and psoriasis lesions after topical application of 5-aminolevulinic acid. *J. Invest. Dermatol.* **110**, 673.
9. Peng, Q., J. Moan, T. Warloe, V. Iani, H. B. Steen, A. Bjørseth and J. M. Nesland (1996) Build-up of esterified aminolevulinic acid-derivative-induced porphyrin fluorescence in normal mouse skin. *J. Photochem. Photobiol. B Biol.* **34**, 95-96.
10. Martin, A., W. D. Tope, J. M. Grevelink, J. C. Starr, J. L. Fewkes, T. J. Flotte, T. F. Deutsch and R. R. Anderson (1995) Lack of selectivity of protoporphyrin IX fluorescence for basal cell carcinoma after topical application of 5-aminolevulinic acid: implications for photodynamic therapy. *Arch. Dermatol. Res.* **287**, 665-674.
11. Szeimies, R. M., T. Sassay and M. Landthaler (1994) Penetration potency of topical applied delta aminolevulinic acid for photodynamic therapy of basal cell carcinoma. *Photochem. Photobiol.* **59**, 73-76.
12. Peng, Q., T. Warloe, J. Moan, H. Heyerdahl, H. B. Steen, J. M. Nesland and K.-E. Giercksky (1995) Distribution of 5-aminolevulinic acid-induced porphyrins in noduloulcerative basal cell carcinoma. *Photochem. Photobiol.* **62**, 906-913.
13. Fritsch, C., C. Abels, G. Goetz, W. Stahl, K. Bolsen, T. Ruzicka, G. Goerz and H. Sies (1997) Porphyrins preferentially accumulate in a melanoma following intravenous injection of 5-aminolevulinic acid. *Biol. Chem.* **378**, 51-57.
14. Lowry, O. H., N. J. Rosenbrough, A. L. Farr and R. J. Randall (1951) Protein measurement with the folin phenol reagent. *J. Biol. Chem.* **193**, 265-275.
15. Seubert, A. and S. Seubert (1982) High-performance liquid chromatographic analysis of porphyrins and their isomers with radial compression columns. *Anal. Biochem.* **124**, 303-307.
16. Fritsch, C., J. Batz, K. Bolsen, K. W. Schulte, M. Zumdick, T. Ruzicka and G. Goerz (1997) *Ex vivo* application of ALA induces high and specific porphyrin levels in human skin tumors. Possible basis for selective photodynamic therapy. *Photochem. Photobiol.* **66**, 114-118.
17. Warloe, T., Q. Peng, H. B. Steen and K. E. Giercksky (1992) Localization of porphyrins in human basal cell carcinoma and normal skin tissue induced by topical application of 5-aminolevulinic acid. In *Photodynamic Therapy and Biomedical Lasers*. (Edited by P. Spinelli, M. Del Fante and R. Marchesini), pp. 454-458. Elsevier, Amsterdam.
18. Stringer, M. R., P. Collins, D. J. Robinson, G. I. Stables and R. A. Sheehan-Dare (1996) The accumulation of protoporphyrin IX in plaque psoriasis after topical application of 5-aminolevulinic acid indicates a potential for photodynamic therapy. *J. Invest. Dermatol.* **107**, 76-81.
19. Malik, Z., G. Kostenich, L. Roitman, B. Ehrenberg and A. Orenstein (1995) Topical application of 5-aminolevulinic acid, DMSO and EDTA: protoporphyrin IX accumulation in skin and tumours of mice. *J. Photochem. Photobiol. B Biol.* **28**, 213-218.
20. Gaullier, J. M., K. Berg, Q. Peng, H. Anholt, P. K. Selbo, L. W. Ma and J. Moan (1997) Use of 5-aminolevulinic acid esters to improve photodynamic therapy on cells in culture. *Cancer Res.* **57**, 1481-1486.
21. Kriegmair, M., R. Baumgartner, R. Knüchel, H. Stepp, F. Hofstädter and A. Hofstetter (1996) Detection of early bladder cancer by 5-aminolevulinic acid induced porphyrin fluorescence. *J. Urol.* **155**, 105-109.
22. Fritsch, C., K. Kalka, P. Lehmann, K. W. Schulte, W. Neuse, T. Ruzicka and G. Goerz (1997) Fluorescence detection of 5-aminolevulinic acid-induced porphyrins (photodynamic diagnosis) in dermatology. *Z. Hautkr. H+G* **8**, 570-576.